

Refer to: Rosove MH: Agranulocytosis and antithyroid drugs.
West J Med 126:339-343, May 1977

Agranulocytosis and Antithyroid Drugs

MICHAEL H. ROSOVE, MD, Santa Monica

Propylthiouracil and methimazole are used widely in the treatment of hyperthyroid disorders. The most important complication of the use of these drugs is depression of the neutrophilic granulocyte count. Granulocytopenia occurs in about 4 percent and agranulocytosis occurs in about 0.3 percent of treated patients. Although this depression of the granulocyte count is reversible after the drug is discontinued, serious infection frequently accompanies agranulocytosis and accounts for almost all deaths related to the drugs. It is important to be aware of the clinical features of granulocytopenic reactions due to antithyroid drugs.

THE TYPES OF TREATMENT used for the various forms of hyperthyroidism include surgical operation, radioactive iodine and antithyroid drugs. The thiourea derivatives, among which are included thiouracil, methylthiouracil, carbimazole, methimazole and propylthiouracil, have, in addition to radioactive iodine, provided the principal mode of nonsurgical treatment for hyperthyroidism since the institution of their use about 30 years ago. The efficacy of propylthiouracil¹⁻⁴ and methimazole⁵⁻⁷ has been well established. These are the two antithyroid agents in widest use in the United States today, largely because of the relatively low incidence of side effects compared with other agents such as thiouracil.⁸⁻¹⁰

In spite of the efficacy and relative safety of propylthiouracil and methimazole, there are adverse reactions. Among the common adverse ef-

fects are skin rashes, urticaria, drug fever and arthralgias.¹¹ Submaxillary gland enlargement and lymphadenopathy may develop.¹² Hypothyroidism may result if the dosage of drug is not reduced when hyperthyroidism comes under control.⁴ Rarely cholestatic jaundice occurs and may be prolonged up to several months.¹³⁻¹⁹ Rare lupus-like syndromes include skin ulcers,²⁰ splenomegaly,^{20,21} migratory polyarthritis,²² pleuritis and pericarditis,^{23,24} periarteritis²⁵ and renal abnormalities.^{11,21} Serologic abnormalities include hyperglobulinemia,^{11,23} positive lupus erythematosus (LE) cell preparations¹¹ and the presence of antinuclear antibodies.^{11,21,22} Rare hematologic complications of therapy are aplastic anemia,²⁶⁻³¹ pure red cell aplasia²⁷ and thrombocytopenia.^{7,14,16,27}

By far the most frequent adverse hematologic effect of antithyroid drugs, however, and the one of most concern for physicians and patients, is granulocytopenia.^{8-16,23,32-45} Most of the common adverse effects have minor significance, and the potentially serious problems of liver involvement

From the Department of Medicine, University of California, Los Angeles, Center for the Health Sciences.

Submitted June 10, 1976.

Reprint requests to: Michael H. Rosove, MD, Department of Medicine, University of California, Los Angeles, Center for the Health Sciences, Los Angeles, CA 90024.

and lupus-like syndromes are very rare. Agranulocytosis is relatively common, however, affecting about one in 300 patients receiving methimazole or propylthiouracil.¹⁰ It is often accompanied by serious infection and accounts for most of the reported drug related deaths. This paper reviews the clinical features of granulocytopenic reactions due to antithyroid drugs.

Incidence

The terms "granulocytopenia" and "agranulocytosis" are not rigidly defined in the literature, but in general they refer to neutrophilic granulocyte counts in the peripheral blood of less than 1,500 per cu mm and 250 per cu mm, respectively.

There is no apparent age predilection for granulocytopenia induced by antithyroid drugs. Children and adults both are affected. Although the case incidence is higher in females, this probably reflects the higher incidence of hyperthyroid conditions in this group and the resultant exposure to these drugs.³⁸

None of the thiourea derivatives have been free of the complication of granulocytopenia. Although thiouracil was an extremely effective antithyroid agent, the primary reason for its discontinuation was the high incidence of granulocytopenic reactions. Van Winkle and co-workers³³ collected reports of 5,745 cases of hyperthyroidism treated with thiouracil and found a 4.4 percent incidence of leukopenia and a 2.5 percent incidence of granulocytopenia. A similar complication rate was noted in the 1,914 thiouracil-treated patients reviewed by Sikkema and associates³⁷ and the 1,091 patients in Moore's review.⁹ Of these 3,005 patients,^{9,37} 20 of 63 patients in whom agranulocytosis developed, died, for an overall 0.67 percent chance of death from thiouracil therapy. Clearly, safer alternatives were necessary.

It was felt that propylthiouracil would be an ideal answer when early trials by Astwood and co-workers¹ and McGavack and associates³ totaling 175 patients failed to show any leukopenic reactions. However, McCullagh and co-workers³⁴ noted a case among their 110 patients, and subsequently Bartels³⁹ published his report of 672 patients treated preoperatively with propylthiouracil among which there were seven instances of granulocytopenia and three cases of agranulocytosis. Beebe and co-workers⁴¹ reported on 1,651 collected cases with 19 instances of granulocytopenia and one case of fatal agranulocytosis. McGavack and Chevalley¹⁰ reviewed the subject of leuko-

penic reactions due to propylthiouracil and methimazole. They found no reactions in patients receiving less than 25 mg of methimazole per day but a 5.4 percent incidence of decreased granulocyte count in those taking 30 mg or more, with one death in about 900 patients. Among the 931 propylthiouracil-treated patients, there was a 2.8 percent incidence of granulocytopenia in those taking less than 150 mg a day, and a 4.5 percent rate in those taking more than 250 mg a day, with one death. The authors concluded that equally effective doses of propylthiouracil and methimazole had a similar rate of complication with a 0.1 percent risk of death due to therapy, and that there was a relationship between dose and risk of granulocytopenia. This relationship was also noted by Wiberg and Nuttall⁴⁵ who noted two cases of agranulocytosis among 25 patients receiving 120 mg methimazole per day.

Clinical Features

The clinical features of agranulocytosis due to antithyroid drugs are similar among the various agents. The onset of symptoms is usually abrupt. The commonest presenting complaints are fever, malaise, gingivitis and sore throat. Those patients with granulocytopenia without agranulocytosis usually remain asymptomatic. The alarming suddenness with which symptoms of agranulocytosis may develop is well known;³⁷ granulocytes may disappear from peripheral blood over two days.³² Leukopenia may be noted before the development of agranulocytosis.^{9,15,36,37} When agranulocytosis develops it generally does so within the first few months of therapy.^{10,13,14,35,38-42} It may occur as early as ten days after therapy is begun¹⁰ or may be delayed for up to four months or more.^{38,40} Moreover, in patients who previously tolerate a drug well agranulocytosis may develop after repeat courses of therapy with the same drug.^{13,36}

The course of symptomatic agranulocytosis may be fulminant with death due to infection occurring within several days.^{14,40,41} In patients who recover following discontinuation of the drug, granulocytes reappear in the peripheral blood after a period varying from a few days to three weeks, and the granulocyte count returns to normal shortly thereafter.^{8,10,35,38,39,42,46} The course of aplastic anemia due to antithyroid drugs may be prolonged.^{29,30}

Among detailed case reports of 25 patients in whom severe drug-related granulocytopenia developed,^{8,9,13-16,35,36,38-43,45} other simultaneously oc-

curing drug-induced side effects were infrequently noted. In one patient urticaria was present,³⁶ in one generalized lymphadenopathy and splenomegaly occurred⁸ and in four jaundice developed.^{14-16,44}

Bone Marrow

There are no unique diagnostic changes in bone marrow in cases of agranulocytosis due to antithyroid drugs.^{12,14,26,29-31,35-37,41-43,45} Except in the rare instances of pancytopenia which are accompanied by an aplastic marrow, only the granulocyte line is affected. The relative number of granulocytes and their precursors may vary considerably from apparent absence to hyperplasia, although a decrease is most common. The marrow morphology is likely to reflect the time during the clinical course at which the specimen is taken.⁴⁶ The distribution of granulocytes is frequently "left-shifted," and the appearance of "maturation arrest" is common. Among the 14 studied cases of thiouracil-induced agranulocytosis collected by Sikkema and associates,³⁷ features compatible with maturation arrest were seen in five. Whether or not the appearance of the marrow in these cases truly represents a maturation arrest is unknown. Excessive utilization of a small pool of granulocyte precursors may produce features of maturation arrest. Moreover, injury incurred at an early stage of granulocyte differentiation may possibly give the appearance of arrest of maturation at a later stage of development.⁴⁷ The stage of granulocyte development at which injury is induced by the thiourea derivatives is not known. In a single case⁴¹ a peroxidase stain showed no positive staining cells, suggesting that the defect was imposed at the myeloblast stage or earlier. Morphologic changes have been noted in mature granulocytes^{36,43} consisting of cytoplasmic swelling with indistinct margins, and irregularly staining, pyknotic nuclei. The significance of these latter findings is unclear.

Pathogenesis

Information available that might shed some light on the pathogenesis of the granulocytopenia and agranulocytosis related to antithyroid drugs is scanty, and the fundamental questions regarding its mechanism are unanswered. The effect of antithyroid agents on thyroid hormone metabolism has been intensively studied. The most important antithyroid actions are to inhibit the oxidation of iodide to the iodinating species necessary in the

synthesis of iodotyrosines and to inhibit the formation of iodothyronines from the coupling of iodotyrosines.⁴⁸ Much less is known about the extrathyroidal effects, and specifically the effect on the bone marrow.

Thiouracil readily gains entry into marrow tissue. Its level builds up slowly and eventually exceeds concentrations in the thyroid.⁴⁹ In patients with nonthyroidal conditions who were given large doses of thiouracil, high concentrations were found in the marrow.⁵⁰ Drug levels within leukocytes may be several times that of plasma.⁴⁹

Oxygen and glucose utilization by leukocytes is affected. Granulocytes from propylthiouracil-treated rats are found to be hypermetabolic with increased oxygen and glucose utilization especially during phagocytosis, an effect independent of thyroid status.⁵¹ However, control leukocytes incubated *in vitro* with propylthiouracil do not show any change in glucose utilization through a wide range of propylthiouracil concentrations.⁵¹ An apparently conflicting observation is that it is possible to show a small but significant decrease in oxygen consumption in granulocytes and to a lesser extent in erythroid cells in the presence of high concentrations of thiouracil.⁵²

Very few specific enzymes have been studied in relation to the thiourea derivatives. The cytochrome oxidase activity of thyroid tissue is inhibited by 0.002 mol of thiouracil *in vitro*; however, concentrations up to 0.01 mol do not affect bone marrow⁵³ although higher concentrations may be inhibitory, raising the question of the significance of the dosage-time factor in some metabolic systems.³⁷ Leukocytes treated with propylthiouracil show slight decreases in the activity of soluble nicotinamide adenine dinucleotide, reduced form (NADH) oxidase, granule bound nicotinamide adenine dinucleotide phosphate, reduced form (NADPH) oxidase and NADH lactic dehydrogenase.⁵¹ The significance of the preceding observations regarding the development of agranulocytosis is unknown.

Some observations regarding the effect of these agents on nucleic acid metabolism have been made. Thiouracil interferes with ribonucleic acid (RNA) synthesis in slices of rat liver and thymus.⁵⁴ Most interesting is that uracil can reverse the inhibition of reticulocyte maturation by thiouracil *in vitro*, suggesting thiouracil is an antimetabolite which may be overcome by adequate concentrations of a necessary substrate.⁵⁵ Seemingly contrary to the above is the report that orotic acid

incorporation into RNA of young leukocytes in propylthiouracil-treated rats is increased.⁵¹ However, the same increase is noted in thyroidectomized animals, and therefore this observation may reflect the thyroid state of the animal rather than the effect of propylthiouracil. In one case of marrow aplasia in man due to propylthiouracil,²⁸ there was no effect of the drug on incorporation of formate or orotic acid into deoxyribonucleic acid (DNA) or RNA, or on uptake of leucine into protein in the bone marrow after recovery.

Specific antibodies are rarely shown. In two cases of aplastic anemia,^{28,30} no leukoagglutinins or platelet agglutinins were found. In a patient with propylthiouracil-induced leukopenia, skin ulcers, fever, arthritis, splenomegaly and hyperglobulinemia,²⁰ immunofluorescent studies with antihuman globulin suggested the presence of an antibody directed to granulocyte cytoplasm. Serum from the patient agglutinated donor granulocytes, and the agglutination was less pronounced with previous incubation of the serum with propylthiouracil. It was suggested that in this patient there was a hypersensitivity disorder with autoimmune leukopenia related to propylthiouracil administration. However, in most cases of granulocytopenia associated with thiourea derivatives, no clinical features suggesting a hypersensitivity disorder are seen. It is likely that agranulocytosis due to antithyroid drugs is not antibody mediated.

Any proposed mechanism relating granulocytopenic reactions to antithyroid drugs will have to take into consideration some clinical observations. First, in only a small percentage of exposed persons does granulocytopenia develop, and even in fewer is there the severe degree of agranulocytosis. Second, the onset of granulocytopenia is frequently delayed. Third, the incidence increases with higher doses. And last, the granulocyte line is usually the only one affected. At present, it seems probable, as previously suggested,²⁸ that in some persons there is a biochemical susceptibility involving some essential metabolic pathway.

Prevention and Treatment

Unfortunately, there is no way to predict in which patient drug-associated agranulocytosis will develop. It is not clear that following blood counts aids in management since agranulocytosis may develop swiftly. Hence it is most important that patients be warned to discontinue medication and contact their physicians if fever, gingivitis or sore throat develops. Agranulocytosis is usually not

presaged by other drug related side effects such as rash, urticaria or arthralgias. Conversely, there is no evidence that patients in whom side effects do develop have an increased risk of developing agranulocytosis.

The proper approach to patients in whom mild to moderate granulocytopenia develops is not clear. In some patients no further depression of the granulocyte count may ever occur and there may be resolution of granulocytopenia despite continued therapy. Amrhein and co-workers¹¹ reported several patients with propylthiouracil-induced granulocytopenia, and the leukopenia resolved with continued treatment. In half the patients with thiouracil-induced leukopenia reported by Van Winkle and associates,³³ the leukocyte count returned to normal levels in spite of continued therapy. A patient with agranulocytosis due to thiouracil was subsequently given the drug again without complication.³² Granulocytopenia may be associated with hyperthyroidism, and in these instances the granulocyte count may rise with antithyroid therapy.⁴⁵

Very little information is available concerning the safety of changing therapy to another thiourea derivative in patients with granulocytopenic reactions. In a few patients in whom granulocytopenia developed with propylthiouracil administration, there was no improvement in the leukocyte count with thiouracil or methimazole therapy.^{11,38} Two patients who had propylthiouracil-induced agranulocytosis were subsequently given thiouracil. In one, agranulocytosis developed; in the other it did not.³⁹ Patients may tolerate one or more drugs well and yet a severe reaction to another may develop. A patient with methimazole-induced aplastic anemia had had previous exposure to propylthiouracil with only an urticarial reaction.²⁶ Another patient was treated with thiouracil for 3 months and with methylthiouracil for 21 months, and then fatal marrow aplasia developed after carbimazole was given for 4 months.²⁷ Unfortunately, there are inadequate numbers of reported cases and no controlled studies concerning the safety of continued therapy with antithyroid drugs in the presence of granulocytopenia. It would seem wise, however, to be certain of the indication for continued medication, to reduce the dose to the minimum necessary to control hyperthyroidism and to follow blood counts in patients who have significant granulocytopenia.

Once agranulocytosis with overt infection is established, treatment consists of immediate ad-

AGRANULOCYTOSIS AND ANTITHYROID DRUGS

mission to hospital, discontinuation of the drug, specific antimicrobial therapy, treatment of hyperthyroidism and general supportive measures. Isolation techniques and granulocyte transfusions may be beneficial. In survivors, the granulocyte count will usually return to normal levels in a period varying from a few days to three weeks.

REFERENCES

1. Astwood EB, Van der Laan WP: Treatment of hyperthyroidism with propylthiouracil. *Ann Intern Med* 25:813-821, 1946
2. McCullagh EP, Hibbs RE, Schneider RW: Propylthiouracil in treatment of hyperthyroidism. *Am J Med Sci* 214:545-552, 1947
3. McGavack TH, Gerl AJ, Vogel M, et al: A clinical comparison of the effectiveness of 6-n-propylthiouracil and 2-thiouracil as antithyrototoxic agents. *Am J Med* 2:144-149, 1947
4. Solomon DH, Beck JC, Van der Laan WP, et al: Prognosis of hyperthyroidism treated by antithyroid drugs. *JAMA* 152:201-205, 1953
5. Stanley MM, Astwood EB: 1-Methyl-2-mercaptoimidazole: Antithyroid compound highly active in man. *Endocrinology* 44:588-589, 1949
6. Reveno WS, Rosenbaum H: Treatment of hyperthyroidism with 1-Methyl-2-mercaptoimidazole. *JAMA* 143:1407, 1950
7. Stone NW, Petit DW, Starr P: Therapeutic studies in hyperthyroidism; methimazole. *J Clin Endocrin* 12:841-845, 1952
8. Fishberg EH, Vorzimer J: Extrathyroid effects of thiouracil therapy. *JAMA* 128:915-921, 1945
9. Moore FD: Toxic manifestations of thiouracil therapy. *JAMA* 130:315-319, 1946
10. McGavack TH, Chevalley J: Untoward hematologic responses to the antithyroid compounds. *Am J Med* 17:36-40, 1954
11. Amrhein JA, Kenny FM, Ross D: Granulocytopenia, lupus-like syndrome, and other complications of propylthiouracil therapy. *J Pediatr* 76:54-63, 1970
12. Limarzi LR, Ricewasser H: Thiouracil granulocytopenia and agranulocytosis. *Am J Clin Path* 16:306-318, 1946
13. Livingston HJ, Livingston SF: Agranulocytosis and hepatocellular jaundice—Toxic reactions following propylthiouracil therapy. *JAMA* 135:422-425, 1947
14. Colwell AR Jr, Sando DE, Lang SJ: Propylthiouracil-induced agranulocytosis, toxic hepatitis, and death. *JAMA* 148:639-641, 1952
15. Rosenbaum H, Reveno WS: Agranulocytosis and toxic hepatitis from methimazole. *JAMA* 152:27, 1953
16. Shipp JC: Jaundice during methimazole (Tapazole) administration. *Ann Intern Med* 42:701-706, 1955
17. Tennenbaum JI, Dreskin OH: Toxic hepatitis during treatment with methimazole (Tapazole): Report of a case with apparent recovery. *Ohio St Med J* 58:306-307, 1962
18. Martinez-Lopez JI, Greenburg SE, Kling RR: Drug-induced hepatic injury during methimazole therapy. *Gastroenterology* 43:84-87, 1962
19. Fischer MG, Nayer HR, Miller A: Methimazole-induced jaundice. *JAMA* 223:1028-1029, 1973
20. Walzer RA, Einbinder J: Immunoleukopenia as an aspect of hypersensitivity to propylthiouracil. *JAMA* 184:743-746, 1963
21. Sammon TJ, Peden VH, Witzleben C, King JP: Disseminated intravascular coagulation complicating propylthiouracil therapy. *Clin Pediatr* 10:739-742, 1971
22. Hung W, August GP: A "collagen-like" syndrome associated with antithyroid therapy. *J Pediatr* 82:852-854, 1973
23. Wing ES, Asper SP: Observations on the use of propylthiouracil in hyperthyroidism. *Bull Hopkins Hosp* 90:201, 1952
24. Best MM, Duncan CH: A lupus-like syndrome following propylthiouracil therapy. *J Kent Med Assoc* 62:47, 1964
25. McCormick RV: Periarteritis occurring during propylthiouracil therapy. *JAMA* 144:1453-1454, 1950
26. Levine B, Rosenberg DV: Aplastic anemia during the treatment of hyperthyroidism with Tapazole. *Ann Intern Med* 41:844-848, 1954
27. Burrell CD: Fatal marrow aplasia after treatment with carbimazole. *Br Med J* 1:1456-1457, 1956
28. Martelo OJ, Katims RB, Yunis AA: Bone marrow aplasia following propylthiouracil therapy. *Arch Intern Med* 120:587-590, 1967
29. Klesse P, Kruskemper HL: Bone marrow damage from mercaptoimidazoles. *Med Klin* 62(8):295-299, 1967
30. Aksoy M, Sakir E: Aplastic anaemia after propylthiouracil. *Lancet* 1:1379, 1968
31. Rosenberg AH: Methimazole induced aplastic anemia. *Conn Med* 34:32-35, 1970
32. Grainger A, Gregson DA, Pemberton HS: Thiouracil in the treatment of thyrotoxicosis. *Br Med J* 2:343-345, 1945
33. Van Winkle W Jr, Hardy SM, Hazel GR, et al: Clinical toxicity of thiouracil, survey of 5,745 cases. *JAMA* 130:343-347, 1946
34. McCullagh EP, Ryan EJ, Schneider R: Propylthiouracil in treatment of hyperthyroidism. *Cleveland Clin Quart* 13:232, 1946
35. Lozinski E, Siminovich J: Thiouracil in treatment of thyrotoxicosis. *Canad Med Assoc J* 51:422, 1944
36. Gargill SL, Lesses MF: Toxic reactions to thiouracil—Report of cases with one fatality. *JAMA* 127:890-898, 1945
37. Sikkema SH, Thewlis EW, Meyer OO: Sternal marrow studies in thyrotoxicosis treated with thiouracil and review of literature regarding thiouracil effects on blood. *Blood* 1:411-425, 1946
38. Eisenmenger WJ, Steele JM: Leukopenia following the use of propylthiouracil. *JAMA* 135:510-511, 1947
39. Bartels EC: Agranulocytosis during propylthiouracil therapy. *Am J Med* 5:48-52, 1948
40. Juliar B, Harris TL: Fatal agranulocytosis occurring during propylthiouracil therapy. *JAMA* 139:646-647, 1949
41. Beebe RT, Propp S, McClintock JC, et al: Fatal agranulocytosis during treatment of toxic goiter with propylthiouracil. *Ann Intern Med* 34:1035-1040, 1951
42. Croke AR, Berry JW: Agranulocytosis occurring during methimazole ("Tapazole") therapy. *JAMA* 148:45-46, 1952
43. Specht NW, Boehme EJ: Death due to agranulocytosis induced by methimazole therapy. *JAMA* 149:1010, 1952
44. Burrell CD, Fraser R, Doniach D: Low toxicity of carbimazole. *Br Med J* 1:1453-1456, 1956
45. Wiberg JJ, Nuttall FQ: Methimazole toxicity from high doses. *Ann Intern Med* 77:414-416, 1972
46. Braun K: Bone marrow changes in a case of agranulocytosis. *Acta Med Orient* 3:120-122, 1944
47. Damashek W: Leukopenia and agranulocytosis. *Oxford Medicine* 2(3):848(15), 1944
48. Ingbar SH, Woelber KA: The thyroid gland, *In* Williams RH (Ed): *Textbook of Endocrinology*. Philadelphia, W B Saunders, 1974, p 175
49. Williams RH, Kay GA, Jandorf BJ: Thiouracil: Its absorption, distribution, and excretion. *J Clin Invest* 23:613-627, 1944
50. Palmer V: Hyperthyroidism and thiouracil: Preliminary report. *Bull Sch Med Univ Maryland* 28:125, 1944
51. Reed PW, Tepperman J: A paradoxical effect of propylthiouracil on rat polymorphonuclear leukocyte metabolism. *Am J Phys* 216:231-237, 1969
52. Warren CO: The effect of thiouracil on the respiration of bone marrow and leukocytes in vitro. *Am J Phys* 145:71-76, 1945
53. Paschkis KE, Cantarow A, Rakoff AE, et al: Effect of thiouracil on tissue oxidase. *Science* 102:333-334, 1945
54. Lindsay RH, Nakagawa H, Cohen PP: Thiouracil-pyrimidine relationships in thyroid and other tissue. *Endocrin* 76:728-736, 1965
55. Nizet A: Antagonistic action of uracil, thiouracil and thio-urea on reticulocyte ripening. *Science* 115:290-291, 1952